
Mechanisms of clonal evolution in childhood acute lymphoblastic leukemia.

Journal: Nat Immunol

Publication Year: 2015

Authors: Srividya Swaminathan, Lars Klemm, Eugene Park, Elli Papaemmanuil, Anthony Ford, Soo-Mi Kweon, Daniel Trageser, Brian Hasselfeld, Nadine Henke, Jana Mooster, Huimin Geng, Klaus Schwarz, Scott C Kogan, Rafael Casellas, David G Schatz, Michael R Lieber, Mel F Greaves, Markus Muschen

PubMed link: 25985233

Funding Grants: Dual targeting of tyrosine kinase and BCL6 signaling for leukemia stem cell eradication

Public Summary:

Childhood acute lymphoblastic leukemia (ALL) can often be traced to a pre-leukemic clone carrying a prenatal genetic lesion. Postnatally acquired mutations then drive clonal evolution toward overt leukemia. The enzymes RAG1-RAG2 and AID, which diversify immunoglobulin-encoding genes, are strictly segregated in developing cells during B lymphopoiesis and peripheral mature B cells, respectively. Here we identified small pre-BII cells as a natural subset with increased genetic vulnerability owing to concurrent activation of these enzymes. Consistent with epidemiological findings on childhood ALL etiology, susceptibility to genetic lesions during B lymphopoiesis at the transition from the large pre-BII cell stage to the small pre-BII cell stage was exacerbated by abnormal cytokine signaling and repetitive inflammatory stimuli. We demonstrated that AID and RAG1-RAG2 drove leukemic clonal evolution with repeated exposure to inflammatory stimuli, paralleling chronic infections in childhood.

Scientific Abstract:

Childhood acute lymphoblastic leukemia (ALL) can often be traced to a pre-leukemic clone carrying a prenatal genetic lesion. Postnatally acquired mutations then drive clonal evolution toward overt leukemia. The enzymes RAG1-RAG2 and AID, which diversify immunoglobulin-encoding genes, are strictly segregated in developing cells during B lymphopoiesis and peripheral mature B cells, respectively. Here we identified small pre-BII cells as a natural subset with increased genetic vulnerability owing to concurrent activation of these enzymes. Consistent with epidemiological findings on childhood ALL etiology, susceptibility to genetic lesions during B lymphopoiesis at the transition from the large pre-BII cell stage to the small pre-BII cell stage was exacerbated by abnormal cytokine signaling and repetitive inflammatory stimuli. We demonstrated that AID and RAG1-RAG2 drove leukemic clonal evolution with repeated exposure to inflammatory stimuli, paralleling chronic infections in childhood.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/mechanisms-clonal-evolution-childhood-acute-lymphoblastic-leukemia>